



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

journal homepage: www.elsevier.com/locate/saa



Synthesis of a new β -naphthothiazole monomethine cyanine dye for the detection of DNA in aqueous solution

Reda M. El-Shishtawy*, Abdullah M. Asiri, Salem A. Basaif, T. Rashad Sobahi

Faculty of Science, Chemistry Department, King Abdul-Aziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

ARTICLE INFO

Article history:

Received 21 October 2009

Received in revised form 22 January 2010

Accepted 11 February 2010

Keywords:

Monomethine cyanine dyes

Ethidium bromide

DNA

Naphthothiazole

Fluorescence enhancement

Intercalation

ABSTRACT

Novel monomethine cyanine dye (MC) derived from β -naphthothiazole and benzothiazole has been prepared and characterized by ^1H and ^{13}C NMR, FTIR, ESIMS, elemental analyses, absorption and fluorescence spectroscopy. The dye was conveniently synthesized by the condensation of two sulfate heterocyclic quaternary salts. The interaction between calf thymus DNA (ct-DNA) in tris(hydroxymethyl)aminomethane-HCl (Tris-HCl) aqueous buffer solution and MC has been studied with spectral fluorescence method. The binding constant value has been determined by fluorescence titration of MC with ct-DNA concentrations. The result obtained is consistent with an intercalative binding interaction between MC and ct-DNA. Compared with ethidium bromide (EB), MC showed a huge fluorescence enhancement upon mixing with ct-DNA.

© 2010 Elsevier B.V. All rights reserved.